

## Topic: Solving Quadratics by Factorising

Topic/Skill	Definition/Tips	Example
1. Quadratic	<p>A quadratic expression is of the form</p> $ax^2 + bx + c$ <p>where <math>a, b</math> and <math>c</math> are numbers, <math>a \neq 0</math></p>	<p>Examples of quadratic expressions:</p> $x^2$ $8x^2 - 3x + 7$ <p>Examples of non-quadratic expressions:</p> $2x^3 - 5x^2$ $9x - 1$
2. Factorising Quadratics	<p>When a quadratic expression is in the form <math>x^2 + bx + c</math> find the two numbers that <b>add to give b</b> and <b>multiply to give c</b>.</p>	$x^2 + 7x + 10 = (x + 5)(x + 2)$ <p>(because 5 and 2 add to give 7 and multiply to give 10)</p> $x^2 + 2x - 8 = (x + 4)(x - 2)$ <p>(because +4 and -2 add to give +2 and multiply to give -8)</p>
3. Difference of Two Squares	<p>An expression of the form <math>a^2 - b^2</math> can be factorised to give <math>(a + b)(a - b)</math></p>	$x^2 - 25 = (x + 5)(x - 5)$ $16x^2 - 81 = (4x + 9)(4x - 9)$
4. Solving Quadratics ( $ax^2 = b$ )	<p>Isolate the <math>x^2</math> term and square root both sides.</p> <p>Remember there will be a <b>positive and a negative solution</b>.</p>	$2x^2 = 98$ $x^2 = 49$ $x = \pm 7$
5. Solving Quadratics ( $ax^2 + bx = 0$ )	<p><b>Factorise</b> and then <b>solve = 0</b>.</p>	$x^2 - 3x = 0$ $x(x - 3) = 0$ $x = 0 \text{ or } x = 3$
6. Solving Quadratics by Factorising ( $a = 1$ )	<p><b>Factorise</b> the quadratic in the usual way.</p> <p><b>Solve = 0</b></p> <p>Make sure the equation = 0 before factorising.</p>	<p>Solve <math>x^2 + 3x - 10 = 0</math></p> <p>Factorise: <math>(x + 5)(x - 2) = 0</math></p> $x = -5 \text{ or } x = 2$
7. Factorising Quadratics when $a \neq 1$	<p>When a quadratic is in the form</p> $ax^2 + bx + c$ <ol style="list-style-type: none"> <li>Multiply <math>a</math> by <math>c = ac</math></li> <li>Find two numbers that add to give <math>b</math> and multiply to give <math>ac</math>.</li> <li>Re-write the quadratic, replacing <math>bx</math> with the two numbers you found.</li> <li>Factorise in pairs – you should get the same bracket twice</li> <li>Write your two brackets – one will be the repeated bracket, the other will be made of the factors outside each of the two brackets.</li> </ol>	<p>Factorise <math>6x^2 + 5x - 4</math></p> <ol style="list-style-type: none"> <li><math>6 \times -4 = -24</math></li> <li>Two numbers that add to give +5 and multiply to give -24 are +8 and -3</li> <li><math>6x^2 + 8x - 3x - 4</math></li> <li>Factorise in pairs: <math>2x(3x + 4) - 1(3x + 4)</math></li> <li>Answer = <math>(3x + 4)(2x - 1)</math></li> </ol>



8. Solving  
Quadratics by  
Factorising  
( $a \neq 1$ )

**Factorise** the quadratic in the usual way.

**Solve = 0**

Make sure the equation = 0 before factorising.

$$\text{Solve } 2x^2 + 7x - 4 = 0$$

$$\text{Factorise: } (2x - 1)(x + 4) = 0$$

$$x = \frac{1}{2} \text{ or } x = -4$$

